

Collision Induced Absorption (CIA) references

These reference numbers can be found in the data headers of the HITRAN CIA files, see also the CIA Readme file for further details.

- [1] K. M. Smith, D. A. Newnham, Near-infrared absorption cross sections and integrated absorption intensities of molecular oxygen (O_2 , $\text{O}_2\text{-O}_2$, and $\text{O}_2\text{-N}_2$), *Journal of Geophysical Research Atmospheres* 105 (D6) (2000) 7383–7396. doi:10.1029/1999JD901171
- [2] Y. Baranov, W. Lafferty, G. Fraser, Investigation of collision-induced absorption in the vibrational fundamental bands of O_2 and N_2 at elevated temperatures, *J. Mol. Spectrosc.* 233 (1) (2005) 160 – 163. doi:10.1016/j.jms.2005.06.008
- [3] W. J. Lafferty, A. M. Solodov, A. Weber, W. B. Olson, J.-M. Hartmann, Infrared collision-induced absorption by N_2 near $4.3\ \mu\text{m}$ for atmospheric applications: measurements and empirical modeling, *Applied Optics* 35 (30) (1996) 5911. doi:10.1364/AO.35.005911
- [4] H. Tran, C. Boulet, J.-M. Hartmann, Line mixing and collision-induced absorption by oxygen in the A-band: Laboratory measurements, model, and tools for atmospheric spectra computations, *J. Geophys. Res.* 111 (2006) D15210. doi:10.1029/2005JD006869
- [5] M. Vangvichith, H. Tran, J.-M. Hartmann, Line-mixing and collision induced absorption for $\text{O}_2\text{-CO}_2$ mixtures in the oxygen A-band region, *J. Quant. Spectrosc. Radiat. Transfer* 110 (18) (2009) 2212–2216. doi:10.1016/j.jqsrt.2009.06.002
- [6] M. Abel, L. Frommhold, X. Li, K. L. C. Hunt, Collision-induced absorption by H_2 pairs: From hundreds to thousands of Kelvin, *J. Phys. Chem. A* 115 (25) (2011) 6805–6812. doi:10.1021/jp109441f
- [7] M. Abel, L. Frommhold, X. Li, K. L. C. Hunt, Infrared absorption by collisional $\text{H}_2\text{-He}$ complexes at temperatures up to 9000 K and frequencies from 0 to 20000 cm^{-1} , *J. Chem. Phys.* 136 (4) (2012) 044319. doi:10.1063/1.3676405
- [8] G. D. Greenblatt, J. J. Orlando, J. B. Burkholder, A. R. Ravishankara, Absorption measurements of oxygen between 330 and 1140 nm, *J. Geophys. Res.* 95 (1990) 18577. doi:10.1029/JD095iD11p18577
- [9] C. Hermans, Measurement of absorption cross sections and spectroscopic molecular parameters, <http://spectrolab.aeronomie.be/o2.htm>
- [10] Maté, B. and Lugez, C. and Fraser, G. T. and Lafferty, W. J., Absolute intensities for the O_2 $1.27\ \mu\text{m}$ continuum absorption, *J. Geophys. Res.: Atmospheres* 104 (D23) (1999) 30585–30590. doi:10.1029/1999JD900824

- [11] A. Borysow, L. Frommhold, Collision-induced rototranslational absorption spectra of $\text{N}_2\text{-N}_2$ pairs for temperatures from 50 to 300 K, *Astrophys. J.* 311 (1986) 1043–1057. doi:10.1086/164841
- [12] A. Borysow, L. Frommhold, Theoretical Collision-induced Rototranslational Absorption Spectra for Modeling Titan’s Atmosphere: $\text{H}_2\text{-N}_2$ Pairs, *Astrophysical Journal* 303 (1986) 495. doi:10.1086/164096
- [13] A. Borysow, C. Tang, Far Infrared CIA Spectra of $\text{N}_2\text{-CH}_4$ Pairs for Modeling of Titan’s Atmosphere, *Icarus* 105 (1) (1993) 175–183. doi:10.1006/icar.1993.1117
- [14] J. Borysow, L. Frommhold, G. Birnbaum, Collision-induced Rototranslational Absorption Spectra of $\text{H}_2\text{-He}$ Pairs at Temperatures from 40 to 3000 K, *Astrophysical Journal* 326 (1988) 509. doi:10.1086/166112
- [15] G. S. Orton, M. Gustafsson, M. Burgdorf, V. Meadows, Revised *ab initio* models for $\text{H}_2\text{-H}_2$ collision-induced absorption at low temperatures, *Icarus* 189 (2) (2007) 544–549. doi:10.1016/j.icarus.2007.02.003
- [16] A. Borysow, L. Frommhold, Theoretical Collision-induced Rototranslational Absorption Spectra for the Outer Planets: $\text{H}_2\text{-CH}_4$ Pairs, *Astrophysical Journal* 304 (1986) 849. doi:10.1086/164221
- [17] M. Gustafsson, L. Frommhold, The $\text{H}_2\text{-H}$ infrared absorption bands at temperatures from 1000 K to 2500 K, *Astronomy & Astrophysics* 400 (3) (2003) 1161–1162. doi:10.1051/0004-6361:20030100
- [18] M. Gustafsson, L. Frommhold, Infrared Absorption Spectra of Collisionally Interacting He and H Atoms, *Astrophysical Journal* 546 (2) (2001) 1168–1170. doi:10.1086/318311
- [19] A. Borysow, L. Frommhold, Collision-induced Rototranslational Absorption Spectra of $\text{CH}_4\text{-CH}_4$ Pairs at Temperatures from 50 to 300 K, *Astrophysical Journal* 318 (1987) 940. doi:10.1086/165426
- [20] M. Gruszka, A. Borysow, Roto-translational collision-induced absorption of CO_2 for the atmosphere of Venus at frequencies from 0 to 250 cm^{-1} , at temperatures from 200 to 800 K, *Icarus* 129 (1997) 172. doi:10.1006/icar.1997.5773
- [21] R. E. Samuelson, N. R. Nath, A. Borysow, Gaseous abundances and methane supersaturation in Titan’s troposphere, *Planetary and Space Science* 45 (8) (1997) 959–980. doi:10.1016/S0032-0633(97)00090-1
- [22] B. J. Drouin, D. C. Benner, L. R. Brown, M. J. Cich, T. J. Crawford, V. M. Devi, A. Guillaume, J. T. Hodges, E. J. Mlawer, D. J. Robichaud, F. Oyafuso, V. H. Payne, K. Sung, E. H. Wishnow, S. Yu, Multispectrum analysis of the oxygen A-band, J.

- Quant. Spectrosc. Radiat. Transfer 186 (2017) 118. doi:10.1016/j.jqsrt.2016.03.037
- [23] Y. I. Baranov, W. J. Lafferty, G. T. Fraser, Infrared spectrum of the continuum and dimer absorption in the vicinity of the O₂ vibrational fundamental in O₂/CO₂ mixtures, Journal of Molecular Spectroscopy 228 (2) (2004) 432–440. doi:10.1016/j.jms.2004.04.010
 - [24] J.-M. Hartmann, C. Boulet, G. C. Toon, Collision-induced absorption by N₂ near 2.16 μ m: Calculations, model, and consequences for atmospheric remote sensing, J. Geophys. Res.: Atmos. 122 (2017) 2419. doi:10.1002/2016JD025677
 - [25] R. H. Taylor, A. Borysow, L. Frommhold, Concerning the rototranslational absorption spectra of He–CH₄ pairs, J. Mol. Spectrosc. 129 (1988) 45. doi:10.1016/0022-2852(88)90257-3
 - [26] E. Bar-Ziv, S. Weiss, Translational Spectra Due to Collision-Induced Overlap Moments in Mixtures of He with CO₂, N₂, CH₄, and C₂H₆, J. Chem. Phys. 57 (1972) 34. doi:10.1063/1.1677970
 - [27] T. Karman, M. A. J. Koenis, A. Banerjee, D. H. Parker, I. E. Gordon, A. van der Avoird, W. J. van der Zande, G. C. Groenenboom, O₂–O₂ and O₂–N₂ collision-induced absorption mechanisms unravelled, Nature Chem. 10 (2018) 549. doi:10.1038/s41557-018-0015-x
 - [28] R. Wordsworth, Y. Kalugina, S. Lokshantov, A. Vigasin, B. Ehlmann, J. Head, C. Sanders, H. Wang, Transient reducing greenhouse warming on early Mars, Geophys. Res. Lett. 44 (2017) 665. doi:10.1002/2016GL071766
 - [29] F. R. Spiering, W. J. van der Zande, Collision induced absorption in the a¹ Δ (ν =2) $\leftarrow$ X³ Σ_g^- (ν =0) band of molecular oxygen, Phys. Chem. Chem. Phys. 14 (28) (2012) 9923–9928. doi:10.1039/c2cp40961e
 - [30] F. R. Spiering, M. B. Kiseleva, N. N. Filippov, L. van Kesteren, W. J. van der Zande, Collision-induced absorption in the O₂ B-band region near 670 nm, Phys. Chem. Chem. Phys. 13 (2011) 9616–9621. doi:10.1039/C1CP20403C
 - [31] T. Karman, E. Miliordos, K. L. C. Hunt, G. C. Groenenboom, A. van der Avoird, Quantum mechanical calculation of the collision-induced absorption spectra of N₂–N₂ with anisotropic interactions, J. Chem. Phys. 142 (2015) 084306. doi:10.1063/1.4907917
 - [32] R. Thalman, R. Volkamer, Temperature dependent absorption cross-sections of O₂–O₂ collision pairs between 340 and 630 nm and at atmospherically relevant pressure, Phys. Chem. Chem. Phys. 15 (37) (2013) 15371. doi:10.1039/c3cp50968k

- [33] Estimated from N₂-N₂ data from Refs. [2, 3] using Eq. (7) of Ref. [3], which is derived using data from the following paper:
 - V. Menoux, R. L. Doucen, C. Boulet, A. Roblin, A. M. Bouchardy, Collision-induced absorption in the fundamental band of N₂: Temperature dependence of the absorption for N₂-N₂ and N₂-O₂ pairs., *Applied optics* 32 (3) (1993) 263–268. doi:10.1364/AO.32.000263
- [34] L. N. Fletcher, M. Gustafsson, G. S. Orton, Hydrogen Dimers in Giant-planet Infrared Spectra, *Astrophysical Journal Supplement Series* 235 (1) (2018) 24. doi:10.3847/1538-4365/aaa07a
- [35] J.-M. Hartmann, C. Boulet, D. D. Tran, H. Tran, Y. Baranov, Effect of humidity on the absorption continua of CO₂ and N₂ near 4 μ m: Calculations, comparisons with measurements, and consequences for atmospheric spectra, *Journal of Chemical Physics* 148 (5) (2018) 054304. doi:10.1063/1.5019994
- [36] E. Wishnow, K. Sung, et al., The far-infrared collision-induced spectrum of nitrogen over the temperature range 78–129 K, *J. Quant. Spectrosc. Radiat. Transfer*, in preparation (2018)
- [37] Y. Baranov, A. Vigasin, Collision-induced absorption by CO₂ in the region of ν_1 , $2\nu_2$, *J. Mol. Spectrosc.* 193 (2) (1999) 319 – 325. doi:10.1006/jmsp.1998.7743
- [38] Y. Baranov, G. T. Fraser, W. J. Lafferty, A. Vigasin, Collision-induced absorption in the CO₂ fermi triad for temperatures from 211 K to 296 K, in: C. Camy-Peyret, A. Vigasin (Eds.), *Weakly Interacting Molecular Pairs: Unconventional Absorbers of Radiation in the Atmosphere*, Springer, 2003, pp. 149–158. doi:10.1007/978-94-010-0025-3_12
- [39] Y. I. Baranov, Collision-induced absorption in the region of the $\nu_2+\nu_3$ band of carbon dioxide, *J. Mol. Spectrosc.* 345 (2018) 11 – 16. doi:10.1016/j.jms.2017.11.005
- [40] D. N. Chistikov, A. A. Finenko, S. E. Lokshtanov, S. V. Petrov, A. A. Vigasin, Simulation of collision-induced absorption spectra based on classical trajectories and ab initio potential and induced dipole surfaces. I. Case study of N₂–N₂ rototranslational band, *Journal of Chemical Physics* 151 (19). doi:10.1063/1.5125756
- [41] T. Odintsova, E. Serov, A. Balashov, M. Koshelev, A. Koroleva, A. Simonova, M. Tretyakov, N. Filippov, D. Chistikov, A. Finenko, S. Lokshtanov, S. Petrov, A. Vigasin, CO₂–CO₂ and CO₂–Ar continua at millimeter wavelengths, *Journal of Quantitative Spectroscopy and Radiative Transfer* 258. doi:10.1016/j.jqsrt.2020.107400

- [42] F. Thibault, V. Menoux, R. Le Doucen, L. Rosenmann, J.-M. Hartmann, C. Boulet, Infrared collision-induced absorption by O_2 near $6.4 \mu m$ for atmospheric applications: measurements and empirical modeling, *Applied Optics* 36 (3) (1997) 563. doi:10.1364/ao.36.000563
- [43] J. J. Orlando, G. S. Tyndall, K. E. Nickerson, J. G. Calvert, The temperature dependence of collision-induced absorption by oxygen near $6 \mu m$, *Journal of Geophysical Research: Atmospheres* 96 (D11) (1991) 20 755–20 760. doi:10.1029/91JD02042
- [44] F. Vitali, S. Stefani, G. Piccioni, M. Snels, D. Grassi, D. Biondi, A. Boccaccini, New experimental measurements of the Collision Induced Absorptions of H_2 - H_2 and H_2 -He in the 3600 - 5500 cm^{-1} spectral range from 120 to 500 K , *Journal of Quantitative Spectroscopy and Radiative Transfer* 330 (2025) 109255. doi:10.1016/j.jqsrt.2024.109255
- [45] A. O. Koroleva, S. Kassi, H. Fleurbaey, A. Campargue, The collision-induced absorption of H_2 near $1.20 \mu m$: Subatmospheric measurements and validation tests of calculations, *Journal of Quantitative Spectroscopy and Radiative Transfer* 318 (2024) 108948. doi:10.1016/j.jqsrt.2024.108948
- [46] M. Turbet, C. Boulet, T. Karman, Measurements and semi-empirical calculations of CO_2+CH_4 and CO_2+H_2 collision-induced absorption across a wide range of wavelengths and temperatures. Application for the prediction of early Mars surface temperature, *Icarus* 346 (2020) 113762. doi:10.1016/j.icarus.2020.113762
- [47] W. Fakhardji, C. Boulet, H. Tran, J.-M. Hartmann, Direct calculations of the CH_4+CO_2 far infrared collision-induced absorption, *Journal of Quantitative Spectroscopy and Radiative Transfer* 283 (2022) 108148. doi:10.1016/j.jqsrt.2022.108148
- [48] H. Fleurbaey, D. Mondelain, W. Fakhardji, J. M. Hartmann, A. Campargue, Simultaneous collision-induced transitions in H_2O+CO_2 gas mixtures, *Journal of Quantitative Spectroscopy and Radiative Transfer* 285 (2022) 108162. doi:10.1016/j.jqsrt.2022.108162
- [49] H. Tran, J. M. Hartmann, E. Rambinison, M. Turbet, Collision-induced absorptions by pure CO_2 in the infrared: New measurements in the 1150 – 4500 cm^{-1} spectral range and empirical modeling for applications, *Icarus* 422 (2024) 116265. doi:10.1016/j.icarus.2024.116265
- [50] C. Wei, A. Klingberg, C. L. Strand, R. K. Hanson, Measurement of hydrogen and nitrogen via collision-induced infrared absorption, *International Journal of Hydrogen Energy* 93 (2024) 364–373. doi:10.1016/j.ijhydene.2024.10.318

- [51] E. A. Serov, T. A. Galanina, A. O. Koroleva, D. S. Makarov, I. S. Amerkhanov, M. A. Koshelev, M. Y. Tretyakov, D. N. Chistikov, A. A. Finenko, A. A. Vigasin, Continuum absorption in pure N₂ gas and in its mixture with Ar, *Journal of Quantitative Spectroscopy and Radiative Transfer* 328 (2024) 109172. doi:10.1016/j.jqsrt.2024.109172
- [52] G. Moreau, Étude en température et modélisation de l’absorption induite par collision dans les régions des bandes fondamentales pour les mélanges des gaz N₂–Ar et O₂, Ph.D. thesis, Université de Rennes (1999).
URL <https://theses.fr/1999REN10083>
- [53] Coefficients were derived from measurements carried out at 271, 297 and 332 K by cavity ring down spectroscopy and described in the following two papers:
 - D. Mondelain, S. Kassi, A. Campargue, Accurate Laboratory Measurement of the O₂ Collision-Induced Absorption Band Near 1.27 μm , *Journal of Geophysical Research (Atmospheres)* 124 (1) (2019) 414423. doi:10.1029/2018JD029317
 - S. Kassi, S. Guessoum, J. C. A. Abanto, H. Tran, A. Campargue, D. Mondelain, Temperature Dependence of the Collision-Induced Absorption Band of O₂ Near 1.27 μm , *Journal of Geophysical Research (Atmospheres)* 126 (13) (2021) e2021JD034860. doi:10.1029/2021JD034860
- [54] H. Finkenzeller, R. Volkamer, O₂-O₂ CIA in the gas phase: Cross-section of weak bands, and continuum absorption between 297-500 nm, *Journal of Quantitative Spectroscopy and Radiative Transfer* 279 (2022) 108063. doi:10.1016/j.jqsrt.2021.108063
- [55] A. A. Finenko, B. Bézard, I. E. Gordon, D. N. Chistikov, S. E. Lokshtanov, S. V. Petrov, A. A. Vigasin, Trajectory-based Simulation of Far-infrared Collision-induced Absorption Profiles of CH₄-N₂ for Modeling Titan’s Atmosphere, *Astrophysical Journal Supplement Series* 258 (2) (2022) 33. doi:10.3847/1538-4365/ac36d3
- [56] A. Banerjee, J. Mandon, F. Harren, D. H. Parker, Collision-induced absorption between O₂-CO₂ for the $\text{a}^1\Delta_g$ ($v = 1$) $\leftarrow$ $\text{X}^3\Sigma_g$ ($v = 0$) transition of molecular oxygen at 1060 nm, *Physical Chemistry Chemical Physics (Incorporating Faraday Transactions)* 21 (4) (2019) 1805–1811. doi:10.1039/C8CP06778C
- [57] J. Terragni, E. Kohler, V. Kofman, S. Anderson, G. L. Villanueva, S. Faggi, New Measurements of High-Temperature CO₂-CO₂ Collision-Induced Absorption in the 1100-1800 cm⁻¹ Range, *Journal of Quantitative Spectroscopy and Radiative Transfer* 346 (2025) 109595. doi:10.1016/j.jqsrt.2025.109595
- [58] E. M. Adkins, T. Karman, A. Campargue, D. Mondelain, J. T. Hodges, Parameterized model to approximate theoretical collision-induced absorption band shapes for O₂-O₂ and O₂-N₂, *Journal of Quantitative Spectroscopy and Radiative Transfer* 310 (2023) 108732. doi:10.1016/j.jqsrt.2023.108732